



Instituto de Ciências da Terra
Institute of Earth Sciences

Study and development of a solar radiation predicting algorithm based on ECMWF's forecasts and ANNs

Sara Pereira, Paulo Canhoto, Rui Salgado and Maria João Costa



Grand Renewable Energy 2018
International Conference and Exhibition

June 17(Sunday) – June 22(Friday) 2018, Pacifico Yokohama, Japan

Data



Experimental data:

- Évora (May 13 2015 to November 15 2016): Kipp & Zonen CM6B pyranometer
- Sines (July 21 2015 to November 15 2016): LI-COR Li-200R pyranometer
- Cabo da Roca and Penhas Douradas (July 21 2015 to November 15 2016): Kipp & Zonen CM11 pyranometer, IPMA

S. Pereira, P. Canhoto, and R. Salgado, “Spatial and temporal downscaling of solar global radiation and mean air temperature from weather forecast data - an introductory numerical study and validation,” in *Workshop On Earth Sciences 2016*, 2016, pp. 8–11.

Variable	ECMWF variable	Unit
Longitude	longitude	° East
Latitude	latitude	° North
Time step	step	h
Date	date	Days since 1900-01-01 00:00:00
Visibility	p3020	m
Cloud base height	cbh	m
Total column water	tcw	kg/m ²
Total cloud cover	tcc	0-1
10 metre U wind component	u10	m/s
10 metre V wind component	v10	m/s
2 metre temperature	t2m	K
2 metre dew point temperature	d2m	K
Surface solar radiation downwards	ssrd	J/m ²
Total column ozone	tco3	kg/m ²



Penhas Douradas



Cabo da Roca


Évora
Sines

Data SIO, NOAA, U.S. Navy, NGA, GEBCO
Image Landsat / Copernicus

Google Earth

Data das imagens: 14/12/2015 42°13'55.53"N 1°33'31.28"E elev 0 m altitude de visualização 3372.01 km






Influence of clearness index (K_t)


$$K_t = \frac{\text{Predicted daily GHI}}{\text{Daily Irradiation on top of the atmosphere}}$$

Hourly differences between predicted and measured GHI [J/m^2] as a function of daily K_t

Évora





Sines





Influence of clearness index (K_t) and solar zenith angle

Évora




Sines




Influence of other variables

Évora


Mean Air Temperature






Relative Air Humidity




Total Water Column



Sines



ANN development





Test Results:

- Hourly average improvement of 22kJ/m^2 ;
- Hourly simulated values equal or closer to measurements than forecasts 61.84% of times.

ANN parameters and inputs:

- Architecture: fitting network with 3 layers;
 - Activation function: hyperbolic tangent sigmoid;
 - Training function: Bayesian regulation backpropagation;
 - 20 validation checks;
 - 80% training, 20% validation.
-
- Common inputs: calculated solar radiation on top of the atmosphere, predicted GHI, visibility, cloud base height, total column water, total cloud cover, 10 meter wind velocity, 2 meter temperature, relative air humidity and total column ozone.
-
- Clear sky days ($K_t \geq 0.65$):
 - Additional inputs: GHI in clear sky conditions (ASHRAE), hour, day and month;
 - 58 neurons;
 - Deviation of 0.145 from clearness index.
 - Partially cloudy days ($0.4 \leq K_t < 0.65$):
 - 30 neurons;
 - Deviation of 0.150 from clearness index.
 - Overcast days ($K_t < 0.4$):
 - Additional inputs: GHI in clear sky conditions (ASHRAE);
 - 27 neurons;
 - Deviation of 0.145 from clearness index.

Example



Conclusion

- Influence of atmospheric and Sun-Earth geometry related variables in the differences between predictions and measurements of GHI:
 - Clearness index;
 - Zenith angle;
 - Mean air temperature;
 - Relative air humidity;
 - Total column water.
- Development of a corrective ANN:
 - Average hourly improvement of 22 kJ/m^2 ;
 - Simulations equal or better than forecasts 61.84% of the times.

This algorithm is integrated in a more complete model, a global solar radiation and mean air temperature prediction algorithm that will be used in energy management in medium/low temperature solar thermal systems. It is expected that the integration of the algorithm in these types of models or models of other solar energy conversion units, will allow for a better management of renewable energy conversion and a better design and usage of conventional auxiliary systems.

Thank you!

Acknowledgment

This work was carried out under the contract with the company Warmhole, lda for the development of a solar radiation forecast algorithm. The authors wish to acknowledge ECMWF and IPMA for the provision of data and the funding provided by the European Regional Development Fund, included in the COMPETE 2020 (Operational Program Competitiveness and Internationalization) through the ICT project (UID/GEO/ 04683/2013) with the reference POCI-01-0145-FEDER -007690, DNI-A (ALT20-03-0145-FEDER-000011) and ALOP (ALT20-03-0145-FEDER-000004) projects .